

Lattice Points on the Homogeneous Cone

$$4(x^2 + y^2) - 7xy = 19z^2$$

Dr. P. Jayakumar

Professor of Mathematics, Annai Vailankanni Arts and Science College Thanjavur-613 007, T.N, India

V. Pandian

Assistant Professor of Mathematics, A.V.V.M. Sri Pushpam College, Poondi-613 503, Thanjavur, T.N, India

Abstract – Seven different methods of the non-zero non-negative solutions of homogeneous Diophantine equation $4(x^2+y^2)-7xy=19z^2$ are obtained. Introducing the linear transformation $x=u+v, y=u-v, u \neq v \neq 0$ in $4(x^2+y^2)-7xy=19z^2$, it reduces to $u^2+15v^2=19z^2$. We are solved the above equation through various choices and the seven different methods of solutions which are satisfied it. Some interesting relations among the special numbers and the solutions are exposed. The following notations are used: $t_{m,n}$ = Polynomial number of rank n with sides m , P_n^m = Pyramidal number of rank n with sides m , G_n = Gnomonic number of rank n , SO_n = Stella Octangular number of rank n

2010 Mathematics subject classification: 11D25

Keywords – The Method of Factorization, Integer Solutions, Linear Transformation, Relations and Special Numbers.

I. INTRODUCTION

The number theory is king of Mathematics. In particular, the Diophantine equations have a blend of attracted interesting problems. For a broad review of variety of problems, one may try to see [3-12]. In 2014, Jayakumar. P, Meena. J [7] published a paper in finding the integer solutions of the homogeneous Diophantine equation $x^2 + 7y^2 = 16z^2$. In 2015, Jayakumar.P., Prabha.S [6] have published a paper in finding the integer solutions of the homogeneous Diophantine equation $x^2 + 9y^2 = 26z^2$. Inspired by these, we are observed in this work another interesting seven different non-zero integral solutions of the homogeneous Diophantine equation $4(x^2+y^2)-7xy = 19z^2$. Further, some elegant properties among the special numbers and the solutions are observed.

II. DESCRIPTION OF METHOD

Consider the Diophantine equation

$$4(x^2 + y^2) - 7xy = 19z^2 \quad (1)$$

Take the linear transformations $x = u + v,$

$$y = u - v, u \neq v \neq 0. \quad (2)$$

Using (1) in (2), it gives us

$$u^2 + 15v^2 = 19z^2 \quad (3)$$

If we take $z = z(a, b) = a^2 + 15b^2 = (a + i\sqrt{15}b)$

$$(a - i\sqrt{15}b), \quad (4)$$

where a and b non-zero non-negative different integers, then we solve (1) through dissimilar method of solutions of (1) which are furnished below.

2.1 Method: I

We can write 19 as

$$19 = (2 + \sqrt{15}i)(2 - \sqrt{15}i) \quad (5)$$

Using (4) and (5) in (3) and applying the factorization process, this gives

$$(u + i\sqrt{15}v)(u - i\sqrt{15}v) = (2 + i\sqrt{15})(2 - i\sqrt{15})(a + i\sqrt{15}b)^2(a - i\sqrt{15}b)^2$$

It gives us

$$(u + i\sqrt{15}v) = (2 + i\sqrt{15})(a + i\sqrt{15}b)^2 \quad (6)$$

$$(u - i\sqrt{15}v) = (2 - i\sqrt{15})(a - i\sqrt{15}b)^2 \quad (7)$$

Comparing both sides of (6) or (7), we obtain

$$u = u(a, b) = 2a^2 - 30ab - 30b^2$$

$$v = v(a, b) = a^2 + 4ab - 15b^2$$

In sight of (2), the solutions x, y are found to be

$$x = x(a, b) = 3a^2 - 26ab - 45b^2 \quad (8)$$

$$y = y(a, b) = a^2 - 34ab - 15b^2 \quad (9)$$

Hence (4), (8) and (9) gives us two parametric the non-zero different integral values of (1).

Observations:

- $x(a, a) + y(a, a) + 116t_{4,a} = 0$
- $x(c, 2c^2-1) - 3y(c, 2c^2-1) - 72SO_c = 0$
- $x(c, c(c+1)) - 3y(c, c(c+1)) - 152Pc^5 = 0$
- $x(a, a+1) + y(a, a+1) + 60t_{4,a+1} + 60P_a - 4t_{4,a} = 0$
- $x(c, 1) - y(c, 1) - 2t_{4,c} - G_{4c} + 29 = 0$
- $x(2, a) + G_{26a} + 45t_{4,a} \equiv 1 \pmod{2}$
- $y(c, 2) + G_{34c} - t_{4,c} \equiv -1 \pmod{2}$
- $3y(b, b+1) - x(b, b+1) + 76P_c = 0$
- $y(2, d) - x(2, d) + G_{8c} - 30t_{4,d} \equiv 0 \pmod{5}$
- $y(2, 2c) + 60t_{4,c} + G_{68c} \equiv 1 \pmod{2}$

2.2 Method: II

We also write 19 as

$$19 = (-2 + i\sqrt{15})(-2 - i\sqrt{15}) \quad (10)$$

Using (4) and (10) in (3) and applying the factorization process, we obtain

$$(u + i\sqrt{15}v)(u - i\sqrt{15}v) = [(-2 + i\sqrt{15})(-2 - i\sqrt{15})](a + i\sqrt{15}b)^2(a - i\sqrt{15}b)^2$$

Comparing both sides of above, we are found to be the positive and negative factors as

$$(u + i\sqrt{15}v) = [(-2 + i\sqrt{15})(a + i\sqrt{15}b)^2] \quad (11)$$

$$(u - i\sqrt{15}v) = [(-2 - i\sqrt{15})(a - i\sqrt{15}b)^2] \quad (12)$$

Comparing both sides of (11) or (12), we obtain the real and imaginary parts as

$$u = u(a, b) = [-2a^2 - 30ab + 15b^2]$$

$$v = v(a, b) = [a^2 - 4ab - 15b^2]$$

In true of (2), the values of x, y are given by

$$x = x(a, b) = -a^2 - 34ab + 15b^2 \quad (13)$$

$$y = y(a, b) = -3a^2 - 26ab + 45b^2 \quad (14)$$

Hence (4), (13) and (14) gives us two parametric the non-zero different integral values of (1).

Observations:

- $y(a, 2a) - 125t_{4c} = 0$
- $x(2b, b) + 57t_{4,b} = 0$
- $x(2, a) + y(2, a) - 60t_{4,a} + G_{60c} + 17 = 0$

4. $3x(b, b+1) + y(b, b+1) - 76P_c = 0$
5. $x(a, a) + y(a, a) + 4t_{4a} = 0$
6. $y(a, 1) - x(a, 1) + 2t_{4,a} - G_{4a} = 0$
7. $y(2, c) - x(2, c) - 30t_{4,c} - G_{8c} + 7 = 0$
8. $x(1, a) - 15t_{4,a} + G_{17a} = 0$
9. $x(b, 2b) + 9t_{4,b} = 0$
10. $y(2b, b) + 19t_{4,b} = 0$

2.3 Method: III

$$\text{Take } 19 \text{ as } 19 = \frac{(17+i\sqrt{15})(17-i\sqrt{15})}{16} \quad (15)$$

For this choice, the corresponding values of x and y obtained from (2) are represented by

$$x = x(a, b) = \frac{1}{2} [9a^2 + 2ab - 135b^2] \quad (16)$$

$$y = y(a, b) = [4a^2 - 16ab - 60b^2] \quad (17)$$

Since our intension is to find integer solutions, taking a as $7A$ and b as $7B$ in (4), (16) and (17), the related parametric integer values of (1) are found as

$$x = x(A, B) = 18A^2 + 4AB - 270B^2 \quad (18)$$

$$y = y(A, B) = 16A^2 - 64AB - 240B^2 \quad (19)$$

$$z = z(A, B) = 4A^2 + 60B^2 \quad (20)$$

Hence (18), (19) and (20) gives us two parametric the non-zero different integral values of (1).

Observations:

1. $x(a, 1) - y(a, 1) - 2t_{4,a} - G_{34a} \equiv -1 \pmod{2}$
2. $x(c, c+1) + y(c, c+1) + 510t_{4,c+1} + 60P_c - 34t_{4,c} = 0$
3. $y(c, 2c^2-1) + 4z(a, 2a^2-1) - 64SO_c - 32t_{4,c} = 0$
4. $y(a, a(a+1)) + 4z(a, a(a+1)) + 128P_a^5 - 32t_{4,a} = 0$
5. $y(a, a) - y(a, a) - 40t_{4,a} = 0$
6. $y(a, 1) + G_{32c} - 16t_{4,a} \equiv -1 \pmod{2}$
7. $z(a, a) - 64t_{4,a} = 0$
8. $y(2a, a) - 304t_{4,a} = 0$
9. $y(2, c) - z(2, c) + 300t_{4,c} + G_{64c} \equiv 1 \pmod{2}$
10. $x(c, 2c) + 10547t_{4,c} = 0$

2.4 Method: IV

$$\text{Also take } 19 \text{ as } 19 = \frac{(-17+i\sqrt{15})(-17-i\sqrt{15})}{16} \quad (21)$$

Following the analysis presented as in pattern- III and simplifying the corresponding non-zero different integer solutions of (1) are found as

$$x = x(A, B) = -16A^2 - 64AB + 240B^2 \quad (22)$$

$$y = y(A, B) = -18A^2 + AB + 270B^2 \quad (23)$$

$$z = z(a, b) = 4A^2 + 60B^2 \quad (24)$$

Hence (22), (23) and (24) gives us two parametric the non-zero different integral values of (1).

Observations:

1. $y(b, b+1) + 3z(b, b+1) + 6t_{4,b+1} - 540t_{4,b} - 4P_b = 0$
2. $y(a, 1) + 18t_{4,a} - G_{2a} \equiv 1 \pmod{2}$
3. $x(b, b+1) + 16t_{4,b} + G_{32b} \equiv 1 \pmod{2}$
4. $y(a, 2a) - 1078t_{4,a} = 0$
5. $x(a, a) + y(a, a) - 7504t_{4,a} = 0$
6. $x(b, 2b) - 816t_{4,b} = 0$
7. $x(1, b) - x(1, b) - 30t_{4,b} + G_{34b} + 1 = 0$
8. $y(1, a) - 3z(1, a) - 30t_{4,a} - G_{2a} \equiv -1 \pmod{2}$
9. $z(b, 1) - 4t_{4,b} \equiv 0 \pmod{2}$
10. $x(2, a) - 240t_{4,a} + G_{64} \equiv 0 \pmod{5}$

2.5 Method: V

Consider (3) as $u^2 - 4z^2 = 15(z^2 - v^2)$ and write it in the form of ratio as

$$\frac{u + 2z}{z + u} = 15 \frac{z - u}{u - 2z} = \frac{A}{B}, B \neq 0 \quad (25)$$

(25) is equivalent to the system of equations

$$Bu - 15v + (2B - 15A)z = 0 \quad (26)$$

$$-Au - Bv + (B + 2A)z = 0 \quad (27)$$

By the cross multiplication method, the above equations yields as

$$u = -30A^2 - 30AB + 2B^2 \quad (28)$$

$$v = 15A^2 - 4AB - B^2 \quad (29)$$

$$z = -15A^2 - B^2 \quad (30)$$

In sight of (2), the solutions x, y are found to be

$$x = x(A, B) = -15A^2 - 34AB + B^2 \quad (31)$$

$$y = y(A, B) = -45A^2 - 26AB + 3B^2 \quad (32)$$

$$z = z(A, B) = -15A^2 - B^2 \quad (33)$$

Thus (31), (32) and (34) gives us two parametric the non-zero different integral values of (1).

Observations:

1. $x(a, a+1) + y(a, a+1) + 60t_{4,a} + 60P_a - 4t_{4,a+1} = 0$
2. $x(a, 2a+1) + z(a, 2a^2-1) + 34t_{4,a} + 34SO_a = 0$
3. $y(a, a(a+1)) - 3x(a, a(a+1)) - 152P_a^5 = 0$
4. $x(a, a) - y(a, a) - 20t_{4,a} = 0$
5. $y(1, a) + G_{13a} - 3t_{4,a} \equiv 0 \pmod{2}$
6. $z(b, b) + 16t_{4,a} = 0$
7. $x(a, 2a) + 79t_{4,a} = 0$
8. $y(2b, b) - 131t_{4,b} = 0$
9. $y(2, b) + z(2, b) + G_{26b} - 2t_{4,b} \equiv -1 \pmod{2}$
10. $x(a, 1) - y(a, 1) - G_{4c} - 30t_{4,a} \equiv 0 \pmod{3}$

2.6 Method: VI

$$\text{Consider (3) as } u^2 + 15v^2 = 19z^2 \times 1 \quad (34)$$

$$\text{Write } 1 \text{ as } 1 = \frac{(7+i\sqrt{15})(7-i\sqrt{15})}{64} \quad (35)$$

Using (34) and (35) in (3) and applying the factorization process, we obtain

$$(u + i\sqrt{15}v)(u - i\sqrt{15}v) = [(2 + i\sqrt{15})(-2 - i\sqrt{15})]$$

$$(a + i\sqrt{15}b)^2 (a - i\sqrt{15}b) \frac{(7+i\sqrt{15})(7-i\sqrt{15})}{64}$$

Comparing both sides of above, we are found to be the positive and negative factors as

$$(u + i\sqrt{15}v) = [(2 + i\sqrt{15})(a + i\sqrt{15}b) \frac{(7+i\sqrt{15})}{8}]$$

$$(u - i\sqrt{15}v) = [(2 - i\sqrt{15})(a - i\sqrt{15}b) \frac{(7-i\sqrt{15})}{8}]$$

Comparing both sides of (11) or (12), we obtain the real and imaginary parts as

$$u = \frac{1}{8} [-a^2 - 270ab + 15b^2]$$

$$v = \frac{1}{8} [9a^2 - 2ab - 135b^2]$$

In sight of (2), the solutions x, y are found to be

$$x = [a^2 - 34ab - 15b^2] \quad (36)$$

$$y = \frac{1}{4} [-5a^2 - 134ab + 75b^2] \quad (37)$$

Since our intension is to find integer solutions, taking a as 2A and b as 2B in (4), (36) and (37), the related parametric integer values of (1) are found as

$$x = x(A, B) = 4A^2 - 136AB - 60B^2 \quad (38)$$

$$y = y(A, B) = -5A^2 - 134AB + 75B^2 \quad (39)$$

$$z = z(A, B) = 4A^2 + 60B^2 \quad (40)$$

Thus (38), (39) and (40) gives us two parametric the non-zero different integral values of (1).

Observations:

1. $x(c, 2c^2-1) + z(c, 2c^2-1) + 136S_{0c} - 8t_{4,c} = 0$
2. $x(c, c+1) - z(c, c+1) + 120t_{4,c+1} + 136P_c = 0$
3. $x(c, 1) + y(c, 1) + t_{4,c} + G_{135c} \equiv 0 \pmod{2}$
4. $x(c, c) + y(c, c) + 256t_{4,c} = 0$
5. $x(c, c(c+1)) + z(c, c(c+1)) + 272P_c^5 - 8t_{4,c} = 0$
6. $z(1, c) - 60t_{4,c} - 4 = 0$
7. $y(2, 2c) + G_{268c} - 300t_{4,c} \equiv 0 \pmod{3}$
8. $y(c, 1) + z(c, 1) + t_{4,c} + G_{67c} \equiv 0 \pmod{2}$
9. $x(c, 1) - z(c, 1) + G_{68c} \equiv 0 \pmod{3}$
10. $x(2, c) + y(2, c) + G_{134c} - 75t_{4,c} = 0$

2.7 Method: VII

Also we write 1 as $1 = \frac{(1+i\sqrt{15})(1-i\sqrt{15})}{16} \quad (41)$

The following analysis presented as in pattern - VI and simplifying the corresponding non-zero different integer solutions of (1) are found as

$$x = x(A, B) = -10A^2 - 116AB + 150B^2 \quad (42)$$

$$y = y(A, B) = -16A^2 - 64AB + 240B^2 \quad (43)$$

$$z = z(A, B) = 4A^2 + 60B^2 \quad (44)$$

Observations:

1. $x(2, 2c) + z(2, 2c) - 840t_{4,c} + G_{232c} \equiv 0 \pmod{5}$
2. $y(2c, c) - 48t_{4,c} = 0$
3. $x(c, 1) - z(c, 1) + 14t_{4,c} + G_{58c} \equiv 1 \pmod{2}$
4. $y(c, 1) - z(c, 1) - 20t_{4,c} + G_{32c} \equiv 1 \pmod{2}$
5. $x(2, c) + y(2, c) - 390t_{4,c} + G_{116c} \equiv 0 \pmod{5}$
6. $x(c, c) + y(c, c) - 184t_{4,c} = 0$
7. $y(c, c(c+1)) - 4z(c, c(c+1)) + 32t_{4,c} + 128P_c^5 = 0$
8. $y(c, 2c^2-1) - 4z(c, 2c^2-1) + 32t_{4,c} + 64S_{0c} = 0$
9. $y(c, c+1) + z(c, c+1) + 12t_{4,c} - 300t_{4,c+1} + 64P_c = 0$
10. $x(c, 1) + y(c, 1) + 26t_{4,c} + G_{90c} \equiv 1 \pmod{2}$

III. CONCLUSION

In this work we observed various process of determining infinitely a lot of non-zero seven different integer values to the homogeneous Diophantine equation $4(x^2+y^2)-7xy=19z^3$. One may try to find non-negative integer solutions of the above equations together with their similar observations.

REFERENCES

- [1] Dickson, L.E., History of theory of numbers, Vol.11, Chelsea publishing company, New - York (1952)
- [2] Mordell, L.J., Diophantine equation, Academic press, London (1969) Journal of science and Research, Vol (3) Issue 12, 20-22 (December -14)
- [3] Jayakumar. P, Sangeetha, K "Lattice points on the cone $x^2 + 9y^2 = 50z^2$ " International Journal of Science and Research, Vol (3), Issue 12, 20-22 (December 2014)
- [4] Jayakumar P, Kanaga Dhurga, C," On Quadratic Diophantine equation $x^2 + 16y^2 = 20z^2$ " Galois J. Maths, 1(1) (2014), 17-23.

- [5] Jayakumar. P, Kanaga Dhurga. C, "Lattice points on the cone $x^2 + 9y^2 = 50z^2$ " Diophantus J. Math, 3(2) (2014), 61-71
- [6] Jayakumar. P, Prabha. S " On Ternary Quadratic Diophantine equation $x^2 + 15y^2 = 14z^2$ " Archimedes J. Math., 4(3) (2014), 159-164.
- [7] Jayakumar, P, Meena, J "Integral solutions of the Ternary Quadratic Diophantine equation : $x^2 + 7y^2 = 16z^2$ " International Journal of Science and Technology, Vol.4, Issue 4, 1-4, Dec 2014.
- [8] Jayakumar. P, Shankarakalidoss, G "Lattice points on Homogenous cone $x^2 + 9y^2 = 50z^2$ " International journal of Science and Research, Vol (4), Issue 1, 2053-2055, January - 2015.
- [9] Jayakumar. P, Shankarakalidoss. G "Integral points on the Homogenous cone $x^2 + y^2 = 10z^2$ " International Journal for Scientific Research and Development, Vol (2), Issue 11, 234-235, January -2015
- [10] Jayakumar. P, Prabha. S "Integral points on the cone $x^2 + 25y^2 = 17z^2$ " International Journal of Science and Research Vol(4), Issue 1, 2050-52, January 2015.
- [11] Jayakumar.P, Prabha. S, "Lattice points on the cone $x^2 + 9y^2 = 26z^2$ " "International Journal of Science and Research Vol (4), Issue 1, 2050-2052, Januar-2015
- [12] Jayakumar. P, Sangeetha. K, "Integral solution of the Homogeneous Biquadratic Diophantine equation with six unknowns: $(x^3-y^3)z = (W^2 - P^2)R^4$ " "International Journal of Science and Research, Vol(3), Issue 12, 1021-1023 (December-2014)

AUTHORS' PROFILES



(1) **Dr. P. Jayakumar** received the B. Sc, M.Sc degrees in Mathematics from Madras University in 1980 and 1983 and the M. Phil, Ph.D degrees in Mathematics from Bharathidasan University, Thiruchirappalli in 1988 and 2010. Who is now working as Professor of Mathematics, Annai Vailankanni Arts and Science College Thanjavur-613 007, T.N, India

(2) **V. Pandian** received the B.Sc, M.Sc, and MPhil degrees in Mathematics from Bharathidasan University, Thiruchirappalli in 2002, 2004 and 2006. Who is now working as Assistant Professor of Mathematics, A.V.V.M. Sri Pushpam College (Autonomous), Poondi - 613 503, Thanjavur. T. N, India.